

## V11 Endovenous Ablation

### Further information

Get more information and references at  
[www.aboutmyhealth.org](http://www.aboutmyhealth.org)

You can get information locally from:

- Ealing Hospital main switchboard on 020 8967 5000
- Central Middlesex Hospital main switchboard on 020 8965 5733
- Northwick Park and St Mark's Hospitals main switchboard on 020 8864 3232
- Our website – [www.lnwh.nhs.uk](http://www.lnwh.nhs.uk)

### Patient Advice and Liaison Service (PALS)

PALS is a confidential service for people who would like information, help or advice about the services provided by any of our hospitals. To contact PALS:

- At Northwick Park, St Mark's and Central Middlesex Hospitals on 020 8869 5118 or email PALS at [LNWH-tr.PALS@nhs.net](mailto:LNWH-tr.PALS@nhs.net)

- At Ealing Hospital and community services on Freephone 0800 064 1120 or 020 8967 5221 between 9.30 am and 4 pm or email PALS at [LNWH-tr.ehPALS@nhs.net](mailto:LNWH-tr.ehPALS@nhs.net)

Please note that this service does not provide clinical advice so please contact the relevant department directly to discuss any concerns or queries about your upcoming test, examination or operation.

**For a translation of this leaflet or for an English version in large print, audio or Braille please ask a member of staff or call 0800 783 4372.**

Tell us how useful you found this document at  
[www.patientfeedback.org](http://www.patientfeedback.org)



[www.rcseng.ac.uk](http://www.rcseng.ac.uk)



[www.rcsed.ac.uk](http://www.rcsed.ac.uk)



[www.asgbi.org.uk](http://www.asgbi.org.uk)



[www.pre-op.org](http://www.pre-op.org)

## What are varicose veins?

Varicose veins are enlarged and twisted veins in your leg. They are common, affecting up to 3 in 10 people.

More women than men ask for treatment, with just over 3 in 10 women being affected aged 35 to 70. Varicose veins tend to run in families and are made worse by pregnancy and if you do a lot of standing.

Your surgeon has recommended endovenous ablation (EVA), an operation to treat the varicose veins. However, it is your decision to go ahead with the operation or not.

This document will give you information about the benefits and risks to help you to make an informed decision. If you have any questions that this document does not answer, ask your surgeon or the healthcare team.

## How do varicose veins happen?

Veins carry blood up your leg and back to your heart. When we stand up, our blood has to be pumped 'uphill' against gravity. Our calf muscles act as a pump and the veins contain many one-way valves to help the upward flow. Both legs contain a system of deep veins, which are buried within the muscles of your leg, and a system of superficial veins which run just underneath your skin.

Sometimes weaknesses in the walls of the superficial veins cause them to enlarge.

The valves then fail to work properly and blood can flow in the wrong direction.

The result is a build-up of pressure in the veins, which bulge out as varicose veins (see figure 1).

## What are the benefits of surgery?

Your symptoms should improve. EVA should help prevent the symptoms and complications that varicose veins cause.

If you have EVA under a local anaesthetic, your surgeon may not be able to remove all the varicose veins (risk: 1 in 3). You may need to have further treatment involving injections or small cuts (avulsions). Avulsions can be performed on the same day as EVA or on another day.

EVA will not remove fine thread veins. If you are having surgery purely for cosmetic reasons, you need to ask your surgeon if EVA will help. This will give you realistic expectations about the final result.



Figure 1  
Varicose veins

## Are there any alternatives to EVA?

Support stockings can often help the symptoms caused by varicose veins and reduce the risk of complications that varicose veins can cause. Foam sclerotherapy is a similar technique that uses an injection of chemicals to treat the veins. Varicose veins surgery involves disconnecting and removing the superficial veins from the deep veins, using a cut on your groin or the back of your knee.

Your surgeon will be able to discuss the options with you.

## What will happen if I decide not to have the operation?

The varicose veins are unlikely to get better without treatment. The following problems may arise.

- Unsightly appearance.
- Itching, aching and pain.
- Pigmentation (dark discolouration) of the skin around your ankle.
- Infection in your skin (cellulitis).
- Inflammation (phlebitis).
- Ulcers (or sores), which are unusual but can be caused by some types of varicose veins.
- Bleeding from varicose veins.

Your surgeon may be able to recommend an alternative treatment for you.

## What does the operation involve?

Before the operation, your surgeon may need to mark the veins on your leg. You may have a Doppler ultrasound (or Duplex scan) of your legs. The healthcare team will carry out a number of checks to make sure you have the operation you came in for and on the correct side. You can help by confirming to your surgeon and the healthcare team your name and the operation you are having. The operation is performed under a general anaesthetic, local anaesthetic or spinal anaesthetic. Your surgeon or anaesthetist will discuss the options with you and recommend the best form of anaesthesia for you. You may be given antibiotics during the operation to reduce the risk of infection. The operation usually takes about 45 minutes.

### • Laser ablation

Your surgeon will insert a catheter (tube) into the great or small saphenous vein. The great saphenous vein runs from the inside of your ankle up to your groin, and the small saphenous vein runs up the back of your leg to the bend in your knee. Your surgeon will use an ultrasound scan to guide them as they insert the catheter into the vein. They will pass a laser fibre through the catheter and up to the point where the saphenous and deep veins meet. They will inject a special liquid to squash the walls of the vein onto the catheter. Your surgeon will slowly remove the catheter and laser fibre while laser energy pulses are sent down the fibre. The laser energy causes the saphenous vein to close (see figure 2).

### • Radio-frequency ablation

Your surgeon will insert a radio-frequency ablation catheter into the great or small saphenous vein. They will use an ultrasound scan to guide them as they move the catheter to the point where the saphenous and deep veins meet. They will inject a special liquid to squash the walls of the vein onto the catheter. The catheter has an electrode that heats the walls of the vein using radio-frequency energy. Your surgeon will slowly remove the catheter while radio-frequency energy is sent down the electrode. The radio-frequency energy causes the saphenous vein to close (see figure 2). Your surgeon may need to remove other varicose veins through small cuts, or close some veins using injections.




Figure 2

An endovenous ablation

If you have EVA under a local anaesthetic, your surgeon may not be able to remove all the varicose veins during the operation. Your surgeon will remove the catheter and cover any cuts with a dressing. You may be given a compression bandage or stocking to wear.

## What should I do about my medication?

Let your doctor know about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.

## What can I do to help make the operation a success?

If you smoke, stopping smoking several weeks or more before the operation may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound.

- In the week before the operation, do not shave or wax the area where a cut is likely to be made.
- Try to have a bath or shower either the day before or on the day of the operation.
- Keep warm around the time of the operation. Let the healthcare team know if you feel cold.

### **What complications can happen?**

The healthcare team will try to make the operation as safe as possible but complications can happen. Some of these can be serious and can even cause death. You should ask your doctor if there is anything you do not understand. Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you.

#### **1 Complications of anaesthesia**

Your anaesthetist or surgeon will be able to discuss with you the possible complications of having an anaesthetic.

#### **2 General complications of any operation**

- Pain. The healthcare team will give you medication to control the pain and it is important that you take it as you are told so you can move about as advised.
- Bleeding during or after the operation. It is common for your leg to be bruised. Rarely, you will need a blood transfusion or another operation. You may also feel a lump under your skin. This is caused by bruising and settles within a few weeks.
- Infection of the surgical site (wound) (risk: less than 1 in 100). It is usually safe to shower after two days but you should check with the healthcare team. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. An infection usually settles with antibiotics but you may need another operation.
- Unsightly scarring of your skin. The scarring will be red at first and will gradually fade to a fine white line.

- Blood clot in your leg (deep-vein thrombosis – DVT) (risk: 1 in 100). This can cause pain, swelling or redness in your leg, or the veins near the surface of your leg to appear larger than normal. The healthcare team will assess your risk. They will encourage you to get out of bed soon after the operation and may give you injections, medication, or special stockings to wear. Let the healthcare team know straightaway if you think you might have a DVT. If you have had a deep-vein thrombosis in one of your legs, let your surgeon know. This sometimes means that surgery should not be performed on that leg.
- Blood clot in your lung (pulmonary embolus), if a blood clot moves through your bloodstream to your lungs. Let the healthcare team know straightaway if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. If you are at home, call an ambulance or go immediately to your nearest Emergency department.

#### **3 Specific complications of this operation**

- Superficial thrombophlebitis, where one of the superficial veins becomes inflamed (risk: 1 in 2). This can cause pain, which is usually easily controlled with simple painkillers such as paracetamol.
- Developing a lump under a wound caused by blood collecting (haematoma) (risk: 3 in 100).
- Numbness or a tingling sensation along the line of the vein that has been treated. This usually improves after a few weeks.
- Burns to your skin. This is unusual as your surgeon uses fluid to protect your skin.
- Ablation device cracking or breaking (risk: less than 1 in 1,000). Your surgeon will need to make a cut on your leg to remove the broken device.
- Damage to nerves (risk: less than 1 in 1,000). If the nerve to your foot is damaged (peroneal nerve), you may have difficulty walking. You may also experience numbness or burning in the skin over your shin.
- Damage to arteries (risk: less than 1 in 1,000). You may need another operation.
- Continued varicose veins. It is not usually possible to remove every single varicose vein.

## How soon will I recover?

### • In hospital

After the operation you will be transferred to the recovery area. You should be able to go home the same day. If you do go home the same day, a responsible adult should take you home in a car or taxi and stay with you for at least 24 hours. Be near a telephone in case of an emergency.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

### • Returning to normal activities

Do not drive, operate machinery or do any potentially dangerous activities (this includes cooking) for at least 24 hours and not until you have fully recovered feeling, movement and co-ordination.

If you had a general anaesthetic or sedation, you should also not sign legal documents or drink alcohol for at least 24 hours.

To reduce the risk of a blood clot, make sure you follow carefully the instructions of the healthcare team if you have been given medication or need to wear special stockings.

When you go home you may have bandages on your legs. The healthcare team will arrange for you to have your bandages removed. You may then need to wear support stockings. The healthcare team will discuss this with you.

Once at home, be as active as possible. When you are resting, keep your legs raised on a stool. Try to return to normal activities as soon as possible, unless you are told otherwise. You may take longer to recover, especially if you had further treatment involving small cuts.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive while you are taking painkillers that make you drowsy. Do not drive until you are confident about controlling your vehicle and always check your insurance policy and with your doctor.

### • The future

Most people make a full recovery. Any aching, swelling or discomfort caused by the operation should gradually improve.

If surgery was performed for ulcers, these should gradually heal.

Skin pigmentation will stay but is less likely to get worse.

You will notice that the varicose veins have gone as soon as the support stockings or bandages are removed.

Varicose veins can come back, either in the same place or in other parts of your leg (risk: 1 in 3 in 5 years).

### Summary

Varicose veins are a common problem and can lead to complications if left untreated. Support stockings can help to control symptoms but will not remove the varicose veins.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

**Keep this information leaflet. Use it to help you if you need to talk to a healthcare professional.**

### Acknowledgements

Author: Mr Bruce Braithwaite MChir FRCS

Illustrations: Medical Illustration Copyright © Nucleus Medical Art. All rights reserved. [www.nucleusinc.com](http://www.nucleusinc.com)

**This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.**